

AD-NILPOTENT POSITIVELY-GRADED BOREL MODULE SUBALGEBRAS

TIM RIDENOUR AND ADAM SANDLER

ABSTRACT. In this paper, we study certain ad-nilpotent subalgebras contained in the non-zero graded portion of a simple $\mathbb{Z}_n$ -graded Lie algebra. These subalgebras respect the grading on the Lie algebra and are modules for a Borel subalgebra for the grade-zero Lie subalgebra. We show that semisimple elements in such subalgebras lie in the center of the subalgebra, and we provide a classification of these subalgebras whose weight space decompositions have only non-zero weights.

INTRODUCTION

Let $\mathfrak{g}$ be a simple Lie algebra with a $\mathbb{Z}_n$ -grading. Let $\mathfrak{g}_0$ be the (reductive) Lie subalgebra of $\mathfrak{g}$ in grade zero, and let $\mathfrak{b}_0$ be a Borel subalgebra for $\mathfrak{g}_0$. Let $\mathfrak{g}_+$ be the direct sum of the non-zero graded components of $\mathfrak{g}$. Then, $\mathfrak{g}_+$ is a $\mathfrak{b}_0$ -module. In this paper, we study $\mathfrak{b}_0$ -submodules of $\mathfrak{g}_+$ which are also ad-nilpotent subalgebras of $\mathfrak{g}$ which respect the $\mathbb{Z}_n$ -grading on $\mathfrak{g}$.

These subalgebras are a generalization of ad-nilpotent ideals of a Borel subalgebra of a simple Lie algebra, which have been studied extensively since the publication of [9]. Since then, several papers have been written on the classification and enumeration of ad-nilpotent ideals of Borel subalgebras and various generalizations (see [3], [4], [5], [13], [11], [12], [6], [1] to name a few).

In [10], Panyushev further generalized this problem to consider the so-called abelian $\mathfrak{b}_0$ -stable subalgebras of $\mathfrak{g}_1$ for a $\mathbb{Z}_2$ -graded Lie algebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$. The classification of these subalgebras was further studied in [2] and [7], among others. In particular, in [7], a criterion on the set of $\mathfrak{g}_0$ -weights of $\mathfrak{g}_1$ was given to classify the abelian $\mathfrak{b}_0$ -stable subalgebras using methods similar to those from [6]. In this paper, we generalize the methods of [7] to classify the k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebras for a simple $\mathbb{Z}_n$ -graded Lie algebra $\mathfrak{g}$.

At this point, we note a few complications that arise in the general case, where $\mathfrak{g}$ is $\mathbb{Z}_n$ -graded, which were not present in the classification of the abelian $\mathfrak{b}_0$ -stable subalgebras for $\mathbb{Z}_2$ -graded Lie algebras. First, if $\mathfrak{g}_+$ is the non-zero graded portion of $\mathfrak{g}$, the non-zero weight spaces of $\mathfrak{g}_+$ (considered as a $\mathfrak{g}_0$ -module) may have dimension higher than one. Accordingly, it is possible for a k -nilpotent $\mathfrak{b}_0$ -module subalgebra of $\mathfrak{g}_+$ to fail to respect the grading on $\mathfrak{g}$; in other words, there exist ad-nilpotent $\mathfrak{b}_0$ -module subalgebras $\mathfrak{a}$ of $\mathfrak{g}_+$ such that $\mathfrak{a} \neq \bigoplus_{j=1}^{n-1} \mathfrak{a} \cap \mathfrak{g}_j$ (see Remark 3.2 for example). We avoid this complication by only considering the subalgebras which respect the grading on $\mathfrak{g}$.

Second, if n is not prime, an ad-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra $\mathfrak{i}$ may contain non-zero elements of weight 0. The methods of [7] do not apply when

0 is a weight of $\mathfrak{i}$. We discuss this possibility at the end of section one and the beginning of section two. In particular, Proposition 1.5 can be used to show that $\mathfrak{i}$ is the direct sum of a subalgebra $\mathfrak{i}'$ which has all non-zero weights and a subalgebra of the elements of $\mathfrak{i}$ with weight 0. The subalgebra $\mathfrak{i}'$ may then be classified completely by its set of weights in each $\mathfrak{g}_j$.

Organization. In Section 1 of this paper, we provide some preliminary results. We show that any semisimple element in an ad-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra $\mathfrak{i}$ is contained in the center of $\mathfrak{i}$.

In section two, we provide our main result. We first show that any k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra $\mathfrak{i}$ is the direct sum of a central subalgebra and a subalgebra with non-zero $\mathfrak{h}_0$ -weights. We then provide a complete classification of the k -nilpotent positively-graded $\mathfrak{b}_0$ -module algebras with only non-zero $\mathfrak{h}_0$ -weights.

Let Q_0^+ be the positive root lattice of (the semisimple portion of) $\mathfrak{g}_0$. Let P_0 be the set of integral weights of $\mathfrak{g}_0$, and let $\mathcal{P} = P_0 \times \mathbb{Z}_n$. We define a partial order $\preceq_0$ on $\mathcal{P}$ as follows: $(\nu, i) \preceq_0 (\mu, j)$ if $i = j$ and $\mu - \nu \in Q_0^+$. As usual, an antichain of $\mathcal{P}_0$ is a set of elements which are unrelated in this order.

Assume that $\mathfrak{i}$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra of $\mathfrak{g}$. A result from [10] shows that a weight space corresponding to a non-zero weight of a graded component of a simple Lie algebra $\mathfrak{g}$ is one-dimensional. Any positively-graded $\mathfrak{b}_0$ -module subalgebra with non-zero weights is therefore a direct sum of one-dimensional weight spaces. Thus, $\mathfrak{i}$ determines a set of weights in each graded portion of $\mathfrak{g}$, and we call the collection of all of these weights along with the grade for each weight the set of *graded $\mathfrak{h}_0$ -weights* of $\mathfrak{i}$, $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i})$. Let $\mathcal{A}_i$ be the antichain of minimal elements of $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(i)$. The set $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i})$ consists precisely of the weights in $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{g}_+)$ which are larger than some element in $\mathcal{A}_i$. We give a necessary and sufficient condition for an antichain of non-zero weights in $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{g}_+)$ to define a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra.

Finally, in section three of the paper, we provide full classifications of abelian positively-graded $\mathfrak{b}_0$ -module subalgebras with non-zero weights in small cases. In particular, we provide a full classification for $\mathfrak{g}$ of type A_3 with a $\mathbb{Z}_3$ -grading, up to conjugacy. We also provide a $\mathbb{Z}_4$ -graded example for A_3 with the grading arising from an outer automorphism.

Acknowledgments. The first author is grateful to RJ Dolbin for sharing the results of his thesis and for fruitful discussions. We are grateful to Volodymyr Mazorchuk for bringing [1] to our attention.

1. PRELIMINARIES

We first fix some notation that will be used throughout the paper. We describe the set of complex numbers (resp. integers, non-negative integers, integers modulo n) by $\mathbb{C}$ (resp. $\mathbb{Z}$, $\mathbb{Z}_+$, $\mathbb{Z}_n$). We write $i \in \mathbb{Z}_n$ for any element of $\mathbb{Z}_n$ instead of $\bar{i}$. Let $\mathfrak{g}$ be a Lie algebra over $\mathbb{C}$ with Cartan subalgebra $\mathfrak{h}$. The set of roots (resp., positive roots) for $\mathfrak{g}$ are denoted by R (resp., R^+). Let I be an indexing set for a set of simple roots for $\mathfrak{g}$. Fix a set of Chevalley generators for $\mathfrak{g}$: $\{x_{\pm\alpha} \mid \alpha \in R^+\} \cup \{h_i \mid i \in I\}$. The Killing form on $\mathfrak{g}$ is denoted by $\kappa : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$.

1.1. The following proposition is standard. We supply a proof for the ease of the reader.

Proposition. *Every $\mathbb{Z}_n$ -grading on a Lie algebra corresponds to a Lie algebra automorphism of $\mathfrak{g}$ with order n . Furthermore, every automorphism of $\mathfrak{g}$ with order n gives a $\mathbb{Z}_n$ grading on $\mathfrak{g}$.*

Proof. Let $\mathfrak{g} = \bigoplus_{i=0}^{n-1} \mathfrak{g}_i$ be a $\mathbb{Z}_n$ -grading on $\mathfrak{g}$. Let ζ be a primitive n^{th} root of unity. Define $\sigma : \mathfrak{g} \rightarrow \mathfrak{g}$ by linearly extending the assignment given by

$$\sigma(x) = \zeta^i x, \quad x \in \mathfrak{g}_i.$$

For $x \in \mathfrak{g}_i$ and $y \in \mathfrak{g}_j$, $[x, y] \in \mathfrak{g}_{i+j}$, which gives

$$\sigma([x, y]) = \zeta^{i+j} [x, y].$$

Also,

$$[\sigma(x), \sigma(y)] = [\zeta^i x, \zeta^j y] = \zeta^{i+j} [x, y],$$

so σ is a Lie algebra automorphism on $\mathfrak{g}$.

Now, suppose that $\varphi : \mathfrak{g} \rightarrow \mathfrak{g}$ is an automorphism of order n . For $0 \leq i \leq n-1$, define

$$\mathfrak{g}_i := \{x \in \mathfrak{g} \mid \varphi(x) = \zeta^i x\}.$$

Then, $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_{n-1}$ gives a $\mathbb{Z}_n$ grading on $\mathfrak{g}$. □

1.2. Let $\mathfrak{g}$ be of type X_N , and let σ be a finite-order automorphism of $\mathfrak{g}$. We note the following (Theorem 8.6 from [8]):

Theorem 1. *Let $\mathbf{s} = (s_0, s_1, \dots, s_\ell; r)$ where $s_0, s_1, \dots, s_\ell$ are nonnegative relatively prime integers and $r \in \{1, 2, 3\}$. Set $n = r \sum_{i=0}^\ell a_i s_i$.*

(i) *The relations*

$$\sigma_{\mathbf{s}}(E_j) = e^{2\pi i s_j / n} E_j, \quad (j = 0, 1, \dots, \ell)$$

define (uniquely) an n^{th} order automorphism of $\mathfrak{g}$.

(ii) *Every n^{th} order automorphism of $\mathfrak{g}$ is conjugate to some $\sigma_{\mathbf{s}}$.*

(iii) *Let $\mathbf{s}' = (s'_0, s'_1, \dots, s'_\ell; r')$. Then, $\sigma_{\mathbf{s}}$ is conjugate to $\sigma_{\mathbf{s}'}$ by a $\mathfrak{g}$ -automorphism if and only if $r = r'$ and the sequence $(s_0, \dots, s_\ell)$ can be transformed to $(s'_0, \dots, s'_\ell)$ by an automorphism of the affine Dynkin diagram of type $X_N^{(r)}$.*

1.3. Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_{n-1}$ be the $\mathbb{Z}_n$ -grading on $\mathfrak{g}$. Define

$$\mathfrak{g}_+ := \bigoplus_{i=1}^{n-1} \mathfrak{g}_i.$$

By the previous theorem, we may assume that $\mathfrak{h}_0 = \mathfrak{h} \cap \mathfrak{g}_0$ is a Cartan subalgebra of $\mathfrak{g}_0$. Note that both $\kappa|_{\mathfrak{g}_0} : \mathfrak{g}_0 \otimes \mathfrak{g}_0 \rightarrow \mathbb{C}$ and $\kappa|_{\mathfrak{h}_0} : \mathfrak{h}_0 \otimes \mathfrak{h}_0 \rightarrow \mathbb{C}$ are non-degenerate. We denote the non-degenerate, symmetric invariant bilinear form on $\mathfrak{h}_0^*$ induced by the restriction of the Killing form on $\mathfrak{g}$ to $\mathfrak{h}_0$ by

$$(\ , \)_0 : \mathfrak{h}_0^* \otimes \mathfrak{h}_0^* \rightarrow \mathbb{C}$$

Let $\mathfrak{b}_0 \subset \mathfrak{g}_0$ be a Borel subalgebra of $\mathfrak{g}_0$ that contains $\mathfrak{h}_0$. We will now define our primary objects of interest.

Definition. A subset $\mathfrak{i} \subset \mathfrak{g}_+$ is called a *k-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra* if $[\mathfrak{i}, \mathfrak{i}] \subset \mathfrak{i}$, $[\mathfrak{b}_0, \mathfrak{i}] \subset \mathfrak{i}$, $(\text{ad } \mathfrak{i})^k(\mathfrak{i}) = 0$ and $\mathfrak{i} = \bigoplus_{j=1}^{n-1} (\mathfrak{i} \cap \mathfrak{g}_j)$.

1.4. In the remainder of this section, we consider semisimple elements in $\mathfrak{g}_+$ and state an important preliminary result concerning the semisimple elements inside of *k-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebras*.

Proposition. Suppose that $h \in \mathfrak{g}_j$ is semisimple and $[\mathfrak{g}_0, h] = 0$. Then, h is in the center of the subalgebra $\mathfrak{g}^{(j)} := \bigoplus_m \mathfrak{g}_{mj}$. In particular, if j is a unit in $\mathbb{Z}_n$, $h = 0$.

Proof. The proof proceeds by induction with the base case given by assumption. Suppose that $[\mathfrak{g}_{mj}, h] = 0$. Then,

$$\kappa([\mathfrak{g}_{-(m+1)j}, h], \mathfrak{g}_{mj}) = \kappa(\mathfrak{g}_{-(m+1)j}, [h, \mathfrak{g}_{mj}]) = 0.$$

By the non-degeneracy of the Killing form, $[\mathfrak{g}_{-(m+1)j}, h] = 0$. Furthermore, since the centralizer of h is reductive, we have $[\mathfrak{g}_{(m+1)j}, h] = 0$.

In the case that j is a unit in $\mathbb{Z}_n$, $\mathfrak{g} = \bigoplus_m \mathfrak{g}_{mj}$. Since $\mathfrak{g}$ is simple, the center is 0; hence, $h = 0$. □

1.5.

Proposition. Suppose that $\mathfrak{i}$ is a *k-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra*. If $h \in \mathfrak{i} \cap \mathfrak{h}_j$, then h is in the center of $\mathfrak{g}^{(j)}$ and $[h, \mathfrak{i}] = 0$. Furthermore, if j is a unit in $\mathbb{Z}_n$, then $\mathfrak{i} \cap \mathfrak{h}_j = 0$.

Proof. Suppose that $h \in (\mathfrak{i} \cap \mathfrak{h}_j)$ is not in the center of $\mathfrak{g}^{(j)}$. Find $e \in \mathfrak{b}_0$ such that $[e, h] \neq 0$, which exists since $[\mathfrak{b}_0, h] = 0$ implies $[\mathfrak{g}_0, h] = 0$, which gives $h = 0$ by Proposition 1.4.

Since $\mathfrak{i}$ is $\mathfrak{b}_0$ -stable, $[h, e] \in \mathfrak{i}$. Furthermore, since $h \in \mathfrak{h}$ and $[h, e] \neq 0$,

$$0 \neq (\text{ad } h)^{k+1}(e) = (\text{ad } h)^k([h, e]) \in (\text{ad } \mathfrak{i})^k(\mathfrak{i}),$$

which contradicts the *k-nilpotence* of $\mathfrak{i}$.

A similar argument shows that $[h, \mathfrak{i}] = 0$. □

1.6. If n is prime, Proposition 1.5 shows that any *k-nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra* $\mathfrak{i}$ contains no non-zero semisimple elements. The following gives an example of an abelian positively-graded $\mathfrak{b}_0$ -module subalgebra which contains non-zero semisimple elements.

Example. Suppose that $\mathfrak{g}$ is of type E_6 . We use the enumeration of the nodes of the Dynkin diagram as in [8]:




FIGURE 1. Dynkin diagram for E_6

Consider $\mathbf{s} = (0, 1, 0, 0, 0; 2)$. Then, $\sigma_{\mathbf{s}}$ is a fourth-order automorphism, which gives rise to a $\mathbb{Z}_4$ -grading of $\mathfrak{g}$. Let $h = 2h_1 + h_2 - h_4 - 2h_5$. Direct computation shows that $h \in \mathfrak{g}_2$ and $[\mathfrak{g}_0, h] = 0$, so $\mathfrak{i} = \mathbb{C}h$ is an abelian positively-graded $\mathfrak{b}_0$ -module subalgebra.

2. MAIN THEOREM

In this section, we prove our main theorem, classifying all k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebras. We first need to introduce some more notation.

Define R_0 to be the set of roots (i.e., non-zero $\mathfrak{h}_0$ -weights) of $\mathfrak{g}_0$, and let Δ_0 be a set of simple roots for $\mathfrak{g}_0$. For each non-zero $j \in \mathbb{Z}_n$, define R_j to be the set of $\mathfrak{h}_0$ -weights of $\mathfrak{g}_j$. For each $\mu \in R_j$, let $\mathfrak{g}_j^{(\mu)}$ be the $\mathfrak{h}_0$ weight space of $\mathfrak{g}_j$ with $\mathfrak{h}_0$ -weight μ . Let

$$\Sigma_+ = \{(\mu, j) \mid \mu \in R_j, \mu \neq 0, j \neq 0\}$$

Suppose that $\mathfrak{i}$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra of $\mathfrak{g}_+$. By Proposition 1.5, we know that $\mathfrak{i} = \mathfrak{i}' \oplus \mathfrak{c}$ where $\mathfrak{c} \subset \mathfrak{h} \cap \mathfrak{g}_+$, $[\mathfrak{c}, \mathfrak{i}] = 0$, and $\mathfrak{i}'$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra with $\mathfrak{i}' \cap \mathfrak{g}_j^{(0)} = \{0\}$ for all $j \in \mathbb{Z}_n$.

Recall the following lemma from [10], which is restated using the notation of the previous paragraph:

Lemma. *If $(\mu, j) \in \Sigma_+$, then the dimension of the weight space $\mathfrak{g}_j^{(\mu)}$ is 1.*

2.1. Let Q_0 (resp., Q_0^+) be the $\mathbb{Z}$ -span (resp., $\mathbb{Z}_+$ -span) of Δ_0 , and let P_0 be the set of integral weights for $\mathfrak{h}_0$. Define $\mathcal{P}_0 := P_0 \times \mathbb{Z}_n$. Then, $\mathcal{P}_0$ has an addition defined by

$$(\eta_1, j_1) + (\eta_2, j_2) = (\eta_1 + \eta_2, j_1 + j_2).$$

Define a partial order $\preccurlyeq_0$ on $\mathcal{P}_0$ via

$$(\mu_1, j_1) \preccurlyeq_0 (\mu_2, j_2) \iff j_1 = j_2 \text{ and } \mu_2 - \mu_1 \in Q_0^+.$$

We will need the following proposition to prove the main theorem:

Proposition. *Suppose that $\beta_j \in R_{i_j}$, $j = 1, \dots, \ell$, and*

$$\sum_{j=1}^{\ell} (\beta_j, i_j) \preccurlyeq_0 (\lambda, i_1 + \dots + i_{\ell}) \text{ for some } \lambda \in R_{i_1 + \dots + i_{\ell}}.$$

Then, there exist $\mu_j \in R_{i_j}$ with $(\beta_j, i_j) \preccurlyeq_0 (\mu_j, i_j)$ for all j , $\sum_{j=1}^{\ell} \mu_j \in R_{i_1 + \dots + i_{\ell}}$ and

$$(\mu_1 + \dots + \mu_{\ell}, i_1 + \dots + i_{\ell}) \preccurlyeq_0 (\lambda, i_1 + \dots + i_{\ell}).$$

Proof. The result is trivial if $\beta_1 + \dots + \beta_{\ell} \in R_{i_1 + \dots + i_{\ell}}$.

Assume $\beta_1 + \dots + \beta_{\ell} \notin R_{i_1 + \dots + i_{\ell}}$. Since

$$\sum_{j=1}^{\ell} (\beta_j, i_j) = (\beta_1 + \dots + \beta_{\ell}, i_1 + \dots + i_{\ell}) \preccurlyeq_0 (\lambda, i_1 + \dots + i_{\ell}),$$

we can write

$$\lambda - \sum_{j=1}^{\ell} \beta_j = \sum_{k=1}^t \gamma_k$$

for some (not necessarily distinct) $\gamma_k \in \Delta_0, k = 1, \dots, t$.

We prove the result by induction on t .

Suppose $t = 1$. Consider $(\beta_1 + \dots + \beta_{\ell}, \gamma_1)_0$. If $(\beta_1 + \dots + \beta_{\ell}, \gamma_1)_0 < 0$, then $(\beta_r, \gamma_1) < 0$ for some r . Define $\mu_r = \beta_r + \gamma_1$ and $\mu_j = \beta_j, j \neq r$. Then, $\mu_j \in R_{i_j}$ for all j , $(\beta_j, i_j) \preceq_0 (\mu_j, i_j)$ for all j , and

$$\sum_{j=1}^{\ell} \mu_j = \beta_1 + \dots + \beta_{\ell} + \gamma_1 = \lambda \in R_{i_1 + \dots + i_{\ell}}.$$

If $(\beta_1 + \dots + \beta_{\ell}, \gamma_1)_0 \geq 0$, then $(\beta_1 + \dots + \beta_{\ell} + \gamma_1, \gamma_1)_0 = (\lambda, \gamma_1)_0 > 0$, which gives

$$\lambda - \gamma_1 = \sum_{j=1}^{\ell} \beta_j \in R_{i_1 + \dots + i_{\ell}}.$$

Now, suppose the result is true for all $s < t$. Consider $(\beta_1 + \dots + \beta_{\ell}, \gamma_1 + \dots + \gamma_t)_0$.

If $(\beta_1 + \dots + \beta_{\ell}, \gamma_1 + \dots + \gamma_t)_0 < 0$, then $(\beta_r, \gamma_m) < 0$ for some r, m . Define $\nu_r = \beta_r + \gamma_m$ and $\nu_j = \beta_j, j \neq r$. Then, $\nu_j \in R_{i_j}$ and $(\beta_j, i_j) \preceq_0 (\nu_j, i_j)$ for all j . Furthermore,

$$\sum_{j=1}^{\ell} (\nu_j, i_j) \preceq_0 (\lambda, i_1 + \dots + i_{\ell}).$$

Since $\lambda - \sum_{j=1}^{\ell} \nu_j = \gamma_1 + \dots + \gamma_{m-1} + \gamma_{m+1} + \dots + \gamma_t$,

the result follows by the induction hypothesis.

If $(\beta_1 + \dots + \beta_{\ell}, \gamma_1 + \dots + \gamma_t)_0 \geq 0$, then

$$(\beta_1 + \dots + \beta_{\ell} + \gamma_1 + \dots + \gamma_t, \gamma_1 + \dots + \gamma_t)_0 = (\lambda, \gamma_1 + \dots + \gamma_t)_0 > 0.$$

For some q , $(\lambda, \gamma_q)_0 > 0$, which gives $\lambda - \gamma_q \in R_{i_1 + \dots + i_{\ell}}$. Set $\lambda' = \lambda - \gamma_q$. This

gives $\sum_{j=1}^{\ell} (\beta_j, i_j) \preceq_0 (\lambda', i_1 + \dots + i_{\ell})$, and

$$\lambda' - \sum_{j=1}^{\ell} \beta_j = \gamma_1 + \dots + \gamma_{q-1} + \gamma_{q+1} + \dots + \gamma_t.$$

We now may apply the induction hypothesis to obtain the result. $\square$

2.2. Let $\mathfrak{i}$ be a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra of $\mathfrak{g}$. For each j , $\mathfrak{i}_j := \mathfrak{i} \cap \mathfrak{g}_j$ has an $\mathfrak{h}_0$ -weight decomposition, and we denote the set of $\mathfrak{h}_0$ -weights of $\mathfrak{i}_j$ by $\text{wt}_{\mathfrak{h}_0}(\mathfrak{i}_j)$. Define the set of *graded $\mathfrak{h}_0$ -weights* of $\mathfrak{i}$,

$$\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i}) := \{(\mu, j) \in \mathcal{P} \mid \mu \in \text{wt}_{\mathfrak{h}_0}(\mathfrak{i}_j)\}.$$

If 0 is not a weight of $\mathfrak{i}_j$ for all j , then $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i}) \subset \Sigma_+$. Let $\mathcal{A}_{\mathfrak{i}}$ denote the set of elements in $\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i})$ which are minimal with respect to $\preceq_0$. Then,

$$\text{wt}_{\mathfrak{h}_0}^{\text{gr}}(\mathfrak{i}) = \{(\mu, j) \in \Sigma_+ \mid (\nu, j) \preceq_0 (\mu, j) \text{ for some } (\nu, j) \in \mathcal{A}_{\mathfrak{i}}\}.$$

We say that an antichain $\mathcal{A} \subset \Sigma_+$ generates a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra, $\mathfrak{i}$, if $\mathcal{A} = \mathcal{A}_{\mathfrak{i}}$. Moreover, we will abuse terminology and say that $\mathcal{A}$ is a *k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra antichain* of Σ_+ if $\mathcal{A} \subset \Sigma_+$ generates a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra.

Theorem 2. *An antichain $\mathcal{A} \subset \Sigma_+$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra antichain if and only if the following three properties hold:*

- (i) *For any $(\mu_1, j_1), (\mu_2, j_2), \dots, (\mu_{k+1}, j_{k+1}) \in \mathcal{A}$,*

$$(\mu_1, j_1) + (\mu_2, j_2) + \dots + (\mu_{k+1}, j_{k+1}) \not\leq_0 (\lambda, j_1 + \dots + j_{k+1})$$
for any highest weight λ of $\mathfrak{g}_{j_1+\dots+j_{k+1}}$.
- (ii) *For any $(\mu_1, j_1), (\mu_2, j_2), \dots, (\mu_m, j_m) \in \mathcal{A}$ with $j_1 + \dots + j_m \equiv 0$,*

$$(\mu_1, j_1) + (\mu_2, j_2) + \dots + (\mu_m, j_m) \not\leq_0 (\theta, 0)$$
for any highest root θ for $\mathfrak{g}_0$.
- (iii) *If $(\mu_1, j_1), (\mu_2, j_2) \in \mathcal{A}$ and $(\mu_1, j_1) + (\mu_2, j_2) \leq_0 (\nu, j_1 + j_2)$ for some $(\nu, j_1 + j_2) \in \Sigma_+$, there exists $(\alpha, j_1 + j_2) \in \mathcal{A}$ such that $(\alpha, j_1 + j_2) \leq_0 (\nu, j_1 + j_2)$.*

Proof. Suppose that an antichain $\mathcal{A} \subset \Sigma_+$ satisfies the three conditions. Let $\mathfrak{i}_{\mathcal{A}}$ be the vector subspace of $\mathfrak{g}_+$ spanned by the $\mathfrak{h}_0$ -weight vectors in $\mathfrak{g}_i$ with weight β where $(\alpha, i) \leq_0 (\beta, i)$ for some $(\alpha, i) \in \mathcal{A}$. It is clear that $\mathfrak{i}_{\mathcal{A}}$ is $\mathfrak{b}_0$ -stable. For any $k+1$ weight vectors, $E_{\beta_1}, E_{\beta_2}, \dots, E_{\beta_{k+1}}$ in $\mathfrak{i}_{\mathcal{A}}$, we need to show that

$$[E_{\beta_1}, [E_{\beta_2}, [\dots, [E_{\beta_k}, E_{\beta_{k+1}}] \dots]] = 0.$$

Suppose that $[E_{\beta_1}, [E_{\beta_2}, [\dots, [E_{\beta_k}, E_{\beta_{k+1}}] \dots]] \neq 0$. Then

$$(\beta_1 + \dots + \beta_{k+1}, j_1 + \dots + j_{k+1}) \leq_0 (\lambda, j_1 + \dots + j_{k+1})$$

for some highest weight λ of $\mathfrak{g}_{j_1+\dots+j_{k+1}}$. For each (β_ℓ, j_ℓ) , $1 \leq \ell \leq k+1$, we can choose $(\mu_\ell, j_\ell) \in \mathcal{A}$ such that $(\mu_\ell, j_\ell) \leq_0 (\beta_\ell, j_\ell)$. Then,

$$(\mu_1, j_1) + \dots + (\mu_{k+1}, j_{k+1}) \leq_0 (\beta_1, j_1) + \dots + (\beta_{k+1}, j_{k+1}) \leq_0 (\lambda, j_1 + \dots + j_{k+1})$$

A similar argument shows that $[E_{\eta_1}, [E_{\eta_2}, [\dots, [E_{\eta_{m-1}}, E_{\eta_m}] \dots]] = 0$ for any weight vectors $E_{\eta_\ell} \in \mathfrak{g}_{j_\ell}$ for $\mathfrak{i}_{\mathcal{A}}$ with $j_1 + \dots + j_m \equiv 0$.

It remains show that $\mathfrak{i}_{\mathcal{A}}$ is a Lie subalgebra of $\mathfrak{g}_+$. Suppose that $E_{\nu_1} \in \mathfrak{g}_{j_1}$ and $E_{\nu_2} \in \mathfrak{g}_{j_2}$ are $\mathfrak{h}_0$ -weight vectors in $\mathfrak{i}_{\mathcal{A}}$ with $[E_{\nu_1}, E_{\nu_2}] \neq 0$. Then, $j_1 + j_2 \neq 0$ by the second condition, so $(\nu_1 + \nu_2, j_1 + j_2) \in \Sigma_+$. Choose $(\mu_\ell, j_\ell) \in \mathcal{A}$, $\ell = 1, 2$, such that $(\mu_\ell, j_\ell) \leq_0 (\nu_\ell, j_\ell)$. This gives $(\mu_1, j_1) + (\mu_2, j_2) \leq_0 (\nu_1 + \nu_2, j_1 + j_2)$. Hence, by the third condition, there exists $(\alpha, j_1 + j_2) \in \mathcal{A}$ such that $(\alpha, j_1 + j_2) \leq_0 (\nu_1 + \nu_2, j_1 + j_2)$, which shows that $[E_{\nu_1}, E_{\nu_2}] \in \mathfrak{i}_{\mathcal{A}}$ by construction.

Therefore, $\mathfrak{i}_{\mathcal{A}}$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra.

Now, suppose that $\mathfrak{i} \subset \mathfrak{g}_+$ is a k -nilpotent positively-graded $\mathfrak{b}_0$ -module subalgebra with $0 \notin \text{wt}_{\mathfrak{b}_0}^{\text{gr}}(\mathfrak{i})$. It remains to prove that $\mathcal{A}_{\mathfrak{i}}$ satisfies the given conditions.

The first two conditions follow immediately from Proposition 2.1.

To show (iii) suppose that $(\mu_1, i_1), (\mu_2, i_2) \in \mathcal{A}$ and

$$(\mu_1, i_1) + (\mu_2, i_2) \leq_0 (\nu, i_1 + i_2)$$

for some $(\nu, i_1 + i_2) \in \Sigma_+$. By Proposition 2.1, there exist $\mu'_j \in R_{i_j}$, $j = 1, 2$, such that $(\mu_j, i_j) \leq_0 (\mu'_j, i_j)$, $\mu'_1 + \mu'_2 \in R_{i_1+i_2}$ and $(\mu'_1 + \mu'_2, i_1 + i_2) \leq_0 (\nu, i_1 + i_2)$. Since $\mathfrak{i}$ is $\mathfrak{b}_0$ -stable, it follows that $(\mu'_1 + \mu'_2, i_1 + i_2) \in \text{wt}_{\mathfrak{b}_0}^{\text{gr}}(\mathfrak{i})$, and, by definition, there exists $(\alpha, i_1 + i_2) \in \mathcal{A}_{\mathfrak{i}}$ such that $(\alpha, i_1 + i_2) \leq_0 (\mu'_1 + \mu'_2, i_1 + i_2) \leq_0 (\nu, i_1 + i_2)$. $\square$

2.3. The statement of the theorem is greatly simplified in the case when $k = 1$ as we note in the following corollary. We use the standard term *abelian* in the place of 1-nilpotent.

Corollary. *An antichain $\mathcal{A} \subset \Sigma_+$ is an abelian positively-graded $\mathfrak{b}_0$ -module subalgebra antichain if and only if for any $(\mu_1, j_1), (\mu_2, j_2) \in \mathcal{A}$,*

$$(\mu_1, j_1) + (\mu_2, j_2) \not\leq_0 (\lambda, j_1 + j_2)$$

for any highest weight λ of $\mathfrak{g}_{j_1+j_2}$.

3. EXAMPLES

In this section, we apply our methods to classify all of the abelian positively-graded $\mathfrak{b}_0$ -module subalgebras for certain cases where $\mathfrak{g}$ is of type A_3 . We let $R^+ = \{\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3\}$ be the set of positive roots for $\mathfrak{g}$, and let $\{x_{\pm\alpha} \mid \alpha \in R^+\} \cup \{h_i \mid i = 1, 2, 3\}$ be a set of Chevalley generators for $\mathfrak{g}$.

3.1. $\mathbb{Z}_3$ -gradings on A_3 . Let $\mathfrak{g}$ be of type A_3 with the set of positive roots R^+ and Chevalley generators as above. Suppose that $n = 3$. By Theorem 1, the automorphisms of $\mathfrak{g}$ of order 3, up to conjugacy, are given by σ_s where s is one of the three following possibilities:

Case 1: $s = (1, 1, 1, 0; 1)$

Case 2: $s = (2, 0, 1, 0; 1)$

Case 3: $s = (0, 0, 1, 2; 1)$

3.1.1. Case 1. Let $s = (1, 1, 1, 0; 1)$. Then, $\mathfrak{g}_0$ is a reductive with $[\mathfrak{g}_0, \mathfrak{g}_0]$ of type A_1 and a 2-dimensional center. The following elements form a basis of $\mathfrak{g}_j$ for $j \in \mathbb{Z}_3$:

$\mathfrak{g}_0$: $x_{\alpha_3}, h_1, h_2, h_3, x_{-\alpha_3}$

$\mathfrak{g}_1$: $x_{\alpha_1}; x_{\alpha_2+\alpha_3}, x_{\alpha_2}; x_{-\alpha_1-\alpha_2}, x_{-\alpha_1-\alpha_2-\alpha_3}$

$\mathfrak{g}_2$: $x_{-\alpha_1}; x_{-\alpha_2}, x_{-\alpha_2-\alpha_3}; x_{\alpha_1+\alpha_2+\alpha_3}, x_{\alpha_1+\alpha_2}$

We choose the Borel subalgebra of $\mathfrak{g}_0$ to be the subalgebra $\mathfrak{b}_0$ spanned by the set $\{x_{-\alpha_3}, h_1, h_2, h_3\}$. The set of simple roots for (the semisimple portion of) $\mathfrak{g}_0$ is then $\Delta_0 = \{-\alpha_3\}$. This unique simple root is also the highest root of $\mathfrak{g}_0$. The sets R_1 and R_2 of weights of $\mathfrak{g}_1$ and $\mathfrak{g}_2$, respectively, are given with the following poset structure with respect to $\leq_0$:

$$R_1 : (\alpha_1, 1); \quad (\alpha_2 + \alpha_3, 1) \leq_0 (\alpha_2, 1); \quad (-\alpha_1 - \alpha_2, 1) \leq_0 (-\alpha_1 - \alpha_2 - \alpha_3, 1)$$

$$R_2 : (-\alpha_1, 2); \quad (-\alpha_2, 2) \leq_0 (-\alpha_2 - \alpha_3, 2); \quad (\alpha_1 + \alpha_2 + \alpha_3, 2) \leq_0 (\alpha_1 + \alpha_2, 2)$$

Using Corollary 2.3, we find that there are exactly 30 non-empty abelian positively-graded $\mathfrak{b}_0$ -module subalgebra antichains in Σ_+ , which we list below:

1. $\{(\alpha_1, 1)\}$
2. $\{(\alpha_2, 1)\}$
3. $\{(\alpha_2 + \alpha_3, 1)\}$
4. $\{(-\alpha_1 - \alpha_2 - \alpha_3, 1)\}$
5. $\{(-\alpha_1 - \alpha_2, 1)\}$
6. $\{(-\alpha_1 - \alpha_2 - \alpha_3, 1), (\alpha_2, 1)\}$

7. $\{(-\alpha_1, 2)\}$
8. $\{(-\alpha_2 - \alpha_3, 2)\}$
9. $\{(-\alpha_2, 2)\}$
10. $\{(\alpha_1 + \alpha_2, 2)\}$
11. $\{(\alpha_1 + \alpha_2 + \alpha_3, 2)\}$
12. $\{(\alpha_1 + \alpha_2, 2), (-\alpha_2 - \alpha_3, 2)\}$
13. $\{(\alpha_1, 1), (\alpha_1 + \alpha_2, 2)\}$
14. $\{(\alpha_1, 1), (\alpha_1 + \alpha_2 + \alpha_3, 2)\}$
15. $\{(\alpha_1, 1), (-\alpha_2 - \alpha_3, 2)\}$
16. $\{(\alpha_1, 1), (-\alpha_2, 2)\}$
17. $\{(-\alpha_1, 2), (-\alpha_1 - \alpha_2 - \alpha_3, 1)\}$
18. $\{(-\alpha_1, 2), (-\alpha_1 - \alpha_2, 1)\}$
19. $\{(-\alpha_1, 2), (\alpha_2, 1)\}$
20. $\{(-\alpha_1, 2), (\alpha_2 + \alpha_3, 1)\}$
21. $\{(\alpha_2, 1), (\alpha_1 + \alpha_2, 2)\}$
22. $\{(\alpha_2, 1), (\alpha_1 + \alpha_2 + \alpha_3, 2)\}$
23. $\{(\alpha_2 + \alpha_3, 1), (\alpha_1 + \alpha_2, 2)\}$
24. $\{(\alpha_2 + \alpha_3, 1), (\alpha_1 + \alpha_2 + \alpha_3, 2)\}$
25. $\{(-\alpha_2 - \alpha_3, 2), (-\alpha_1 - \alpha_2 - \alpha_3, 1)\}$
26. $\{(-\alpha_2 - \alpha_3, 2), (-\alpha_1 - \alpha_2, 1)\}$
27. $\{(-\alpha_2, 2), (-\alpha_1 - \alpha_2 - \alpha_3, 1)\}$
28. $\{(-\alpha_2, 2), (-\alpha_1 - \alpha_2, 1)\}$
29. $\{(-\alpha_1 - \alpha_2 - \alpha_3, 1), (\alpha_2, 1), (-\alpha_1, 2)\}$
30. $\{(\alpha_1 + \alpha_2, 2), (-\alpha_2 - \alpha_3, 2), (\alpha_1, 1)\}$


3.1.2. Case 2. Let $\mathbf{s} = (2, 0, 1, 0; 1)$. Then, $\mathfrak{g}_0$ is a reductive algebra with a one-dimensional center, and $[\mathfrak{g}_0, \mathfrak{g}_0]$ is of type $A_1 \times A_1$. The following elements form a basis of $\mathfrak{g}_j$ for $j \in \mathbb{Z}_3$:

$\mathfrak{g}_0$: $x_{\alpha_1}, x_{\alpha_3}, h_1, h_2, h_3, x_{-\alpha_1}, x_{-\alpha_3}$

$\mathfrak{g}_1$: $x_{\alpha_1+\alpha_2+\alpha_3}, x_{\alpha_1+\alpha_2}, x_{\alpha_2+\alpha_3}, x_{\alpha_2}$

$\mathfrak{g}_2$: $x_{-\alpha_2}, x_{-\alpha_1-\alpha_2}, x_{-\alpha_2-\alpha_3}, x_{-\alpha_1-\alpha_2-\alpha_3}$

We set the Borel subalgebra of $\mathfrak{g}_0$ to be the subalgebra $\mathfrak{b}_0$ spanned by the set $\{x_{\alpha_1}, x_{\alpha_3}, h_1, h_2, h_3\}$. The set of simple roots for (the semisimple portion of) $\mathfrak{g}_0$ is then $\Delta_0 = \{\alpha_1, \alpha_3\}$. These two roots are the highest roots of $\mathfrak{g}_0$. The sets R_1 and R_2 of weights of $\mathfrak{g}_1$ and $\mathfrak{g}_2$, respectively, are given with the following poset structure with respect to $\preceq_0$, where an arrow $(\mu, i) \rightarrow (\nu, i)$ indicates that $(\mu, i) \preceq_0 (\nu, i)$:



Using Corollary 2.3, we find that there are exactly 11 non-empty abelian positively-graded $\mathfrak{b}_0$ -module subalgebra antichains in Σ_+ , which we list below:

1. $\{(\alpha_2, 1)\}$
2. $\{(\alpha_1 + \alpha_2, 1)\}$
3. $\{(\alpha_2 + \alpha_3, 1)\}$
4. $\{(\alpha_1 + \alpha_2 + \alpha_3, 1)\}$
5. $\{(-\alpha_1 - \alpha_2 - \alpha_3, 2)\}$
6. $\{(-\alpha_1 - \alpha_2, 2)\}$
7. $\{(-\alpha_2 - \alpha_3, 2)\}$
8. $\{(-\alpha_2, 2)\}$
9. $\{(\alpha_1 + \alpha_2, 1), (\alpha_2 + \alpha_3, 1)\}$
10. $\{(-\alpha_1 - \alpha_2, 2), (-\alpha_2 - \alpha_3, 2)\}$
11. $\{(-\alpha_2, 2), (\alpha_1 + \alpha_2 + \alpha_3, 1)\}$

3.1.3. Case 3. Consider the $\mathbb{Z}_3$ -grading on $\mathfrak{g}$ given by the automorphism $\mathbf{s} = (0, 0, 1, 2; 1)$. Then, $\mathfrak{g}_0$ has a 1-dimensional center and $[\mathfrak{g}_0, \mathfrak{g}_0]$ is of type A_2 . The following elements form a basis of $\mathfrak{g}_j$ for $j \in \mathbb{Z}_3$:

$$\underline{\mathfrak{g}_0}: x_{-\alpha_1-\alpha_2-\alpha_3}, x_{\alpha_1}, x_{-\alpha_2-\alpha_3}, h_1, h_2, h_3, x_{\alpha_1+\alpha_2+\alpha_3}, x_{-\alpha_1} x_{\alpha_2+\alpha_3}$$

$$\underline{\mathfrak{g}_1}: x_{-\alpha_3}, x_{\alpha_1+\alpha_2}, x_{\alpha_2}$$

$$\underline{\mathfrak{g}_2}: x_{-\alpha_2}, x_{-\alpha_1-\alpha_2}, x_{\alpha_3}$$

We set the Borel subalgebra of $\mathfrak{g}_0$ to be the subalgebra $\mathfrak{b}_0$ spanned by the set $\{x_{-\alpha_1-\alpha_2-\alpha_3}, x_{\alpha_1}, x_{-\alpha_2-\alpha_3}, h_1, h_2, h_3\}$. The set of simple roots for (the semisimple portion of) $\mathfrak{g}_0$ is then $\Delta_0 = \{-\alpha_1 - \alpha_2 - \alpha_3, \alpha_1\}$. The highest root of $\mathfrak{g}_0$ is then $\theta_0 = -\alpha_2 - \alpha_3$. The sets R_1 and R_2 of weights of $\mathfrak{g}_1$ and $\mathfrak{g}_2$, respectively, are given with the following poset structure with respect to $\preceq_0$:

$$R_1 : (\alpha_2, 1) \preceq_0 (\alpha_1 + \alpha_2, 1) \preceq_0 (-\alpha_3, 1)$$

$$R_2 : (\alpha_3, 2) \preceq_0 (-\alpha_1 - \alpha_2, 2) \preceq_0 (-\alpha_2, 2)$$

Using Corollary 2.3, we find that there are exactly 6 non-empty abelian positively-graded $\mathfrak{b}_0$ -module subalgebra antichains in Σ_+ , which we list below:

1. $\{(-\alpha_3, 1)\}$
2. $\{(\alpha_1 + \alpha_2, 1)\}$
3. $\{(\alpha_2, 1)\}$
4. $\{(-\alpha_2, 2)\}$
5. $\{(-\alpha_1 - \alpha_2, 2)\}$
6. $\{(\alpha_3, 2)\}$

3.2. An outer automorphism. Let $\mathfrak{g}$ be of type A_3 with the Chevalley generators given at the beginning of the section. Define the following elements of $\mathfrak{g}$:

$$E_0 = x_{-\alpha_1-\alpha_2} - x_{-\alpha_2-\alpha_3}$$

$$\begin{aligned}
E_1 &= x_{\alpha_1} + x_{\alpha_3} \\
E_2 &= x_{\alpha_2} \\
F_0 &= -x_{\alpha_1+\alpha_2} + x_{\alpha_2+\alpha_3} \\
F_1 &= x_{-\alpha_1} + x_{-\alpha_3} \\
F_2 &= x_{-\alpha_2} \\
H_1 &= h_1 + h_3 \\
H_2 &= h_2 \\
E_{\gamma_1+\gamma_2} &= x_{\alpha_1+\alpha_2} + x_{\alpha_2+\alpha_3} \\
E_{2\gamma_1+\gamma_2} &= x_{\alpha_1+\alpha_2+\alpha_3} \\
E_{\gamma_0+\gamma_2} &= x_{-\alpha_1} - x_{-\alpha_3} \\
E_{\gamma_0+\gamma_1+\gamma_2} &= h_1 - h_3 \\
F_{\gamma_1+\gamma_2} &= x_{-\alpha_1-\alpha_2} + x_{-\alpha_2-\alpha_3} \\
F_{2\gamma_1+\gamma_2} &= x_{-\alpha_1-\alpha_2-\alpha_3} \\
F_{\gamma_0+\gamma_2} &= -x_{\alpha_1} + x_{\alpha_3}
\end{aligned}$$

Consider the $\mathbb{Z}_4$ -grading on $\mathfrak{g}$ given by the automorphism $\sigma_{(1,1,0;2)}$. Here, $\mathfrak{h}_0$ is spanned by H_1 and H_2 . The elements $\gamma_\ell \in \mathfrak{h}_0^*$, $\ell = 0, 1, 2$, are given by $[h, E_\ell] = \gamma_\ell(h)E_\ell$ for all $h \in \mathfrak{h}_0$. Note that $\gamma_0 + \gamma_1 + \gamma_2 = 0$ as elements in $\mathfrak{h}_0^*$. For any γ , $[h, E_\gamma] = \gamma(h)E_\gamma$ and $[h, F_\gamma] = -\gamma(h)F_\gamma$ for all $h \in \mathfrak{h}_0$.

Using the above elements, the following form a basis of $\mathfrak{g}_j$ for $j \in \mathbb{Z}_4$:

$$\underline{\mathfrak{g}}_0: E_1, H_1, F_1; H_2$$

$$\underline{\mathfrak{g}}_1: E_0; E_{2\gamma_1+\gamma_2}, E_{\gamma_1+\gamma_2}, E_2$$

$$\underline{\mathfrak{g}}_2: F_{\gamma_0+\gamma_2}, E_{\gamma_0+\gamma_1+\gamma_2}, E_{\gamma_0+\gamma_2}$$

$$\underline{\mathfrak{g}}_3: F_0; F_2, F_{\gamma_1+\gamma_2}, F_{2\gamma_1+\gamma_2}$$

We set the Borel subalgebra of $\mathfrak{g}_0$ to be the subalgebra $\mathfrak{b}_0$ spanned by the set $\{E_1, H_1, H_2\}$. The set of simple roots for (the semisimple portion of) $\mathfrak{g}_0$ is then $\Delta_0 = \{\gamma_1\}$. This simple root is also the highest root for $\mathfrak{g}_0$. The sets R_1 , R_2 and R_3 of weights of $\mathfrak{g}_1$, $\mathfrak{g}_2$ and $\mathfrak{g}_3$, respectively, are given with the following poset structure with respect to $\preceq_0$:

$$\begin{aligned}
R_1: & \quad (\gamma_0, 1); \\
& \quad (\gamma_2, 1) \preceq_0 (\gamma_1 + \gamma_2, 1) \preceq_0 (2\gamma_1 + \gamma_2, 1) \\
R_2: & \quad (\gamma_0 + \gamma_2, 2) \preceq_0 (\gamma_0 + \gamma_1 + \gamma_2, 2) \preceq_0 (-\gamma_0 - \gamma_2, 2) \\
R_3: & \quad (-\gamma_0, 3); \\
& \quad (-2\gamma_1 - \gamma_2, 3) \preceq_0 (-\gamma_1 - \gamma_2, 3) \preceq_0 (-\gamma_2, 3)
\end{aligned}$$

Using Corollary 2.3, we find that there are exactly 20 non-empty abelian positively-graded $\mathfrak{b}_0$ -module subalgebra antichains in Σ_+ , which we list below:

1. $\{(\gamma_0, 1)\}$
2. $\{(2\gamma_1 + \gamma_2, 1)\}$
3. $\{(\gamma_1 + \gamma_2, 1)\}$

4. $\{(\gamma_2, 1)\}$
5. $\{(-\gamma_0 - \gamma_2, 2)\}$
6. $\{(-\gamma_0, 3)\}$
7. $\{(-\gamma_2, 3)\}$
8. $\{(-\gamma_1 - \gamma_2, 3)\}$
9. $\{(-2\gamma_1 - \gamma_2, 3)\}$
10. $\{(2\gamma_1 + \gamma_2, 1), (-\gamma_0 - \gamma_2, 2)\}$
11. $\{(\gamma_1 + \gamma_2, 1), (-\gamma_0 - \gamma_2, 2)\}$
12. $\{(\gamma_0, 1), (-\gamma_2, 3)\}$
13. $\{(\gamma_0, 1), (-\gamma_1 - \gamma_2, 3)\}$
14. $\{(\gamma_0, 1), (-2\gamma_1 - \gamma_2, 3)\}$
15. $\{(2\gamma_1 + \gamma_2, 1), (-\gamma_2, 3)\}$
16. $\{(-\gamma_0, 3), (2\gamma_1 + \gamma_2, 1)\}$
17. $\{(-\gamma_0, 3), (\gamma_1 + \gamma_2, 1)\}$
18. $\{(-\gamma_0, 3), (\gamma_2, 1)\}$
19. $\{(-\gamma_0 + \gamma_2, 2), (-\gamma_2, 3)\}$
20. $\{(-\gamma_0 + \gamma_2, 2), (-\gamma_1 - \gamma_2, 3)\}$

Remark. The elements $E_{\gamma_1 + \gamma_2} \in \mathfrak{g}_1$ and $F_0 \in \mathfrak{g}_3$ have the same $\mathfrak{h}_0$ -weight. Let $v = E_{\gamma_1 + \gamma_2} + F_0$. The $\mathfrak{b}_0$ -submodule of $\mathfrak{g}_+$ generated by v is an abelian subalgebra of $\mathfrak{g}_+$ that is not positively graded.

REFERENCES

- [1] K. Baur and V. Mazorchuk. Combinatorial analogues of ad-nilpotent ideals for untwisted affine Lie algebras. *J. Algebra (to appear)*, 2011. arXiv:1108.3659.
- [2] P. Cellini, P. Frajria, and P. Papi. Abelian Subalgebras in $\mathbb{Z}_2$ -graded Lie Algebras and Affine Weyl Group. *Int. Math. Res. Notices*, 2004(43):2281–2304, 2004.
- [3] P. Cellini and P. Papi. ad-nilpotent ideals of a Borel subalgebra. *J. Algebra*, 225:130–141, 2000.
- [4] P. Cellini and P. Papi. ad-nilpotent ideals of a Borel subalgebra II. *J. Algebra*, 258:112–121, 2002.
- [5] P. Cellini and P. Papi. ad-nilpotent ideals of Borel subalgebras and affine weyl groups. *Adv. Math.*, no. 187, 2004.
- [6] V. Chari, R.J. Dolbin, and T. Ridenour. Ideals in parabolic subalgebras of simple Lie algebras. *Contemporary Mathematics*, (490):47–60, 2009.
- [7] R.J. Dolbin. $\mathbb{Z}_2$ -graded Lie algebras. PhD thesis, University of California, Riverside, 2010.
- [8] V. Kac. *Infinite dimensional Lie algebras*. Cambridge University Press, third edition, 1990.
- [9] B. Kostant. The set of abelian ideals of a Borel subalgebra, Cartan decompositions, and discrete series representation. *Int. Math. Res. Notices*, pages 225–252, 1998.
- [10] D. Panyushev. Isotropy Representations, Eigenvalues of a Casimir element, and commutative Lie subalgebras. *J. London Math Society*, 64(2):61–80, 2001.
- [11] D. Panyushev. Abelian ideals of a Borel subalgebra and long positive roots. *IMRN*, 35, 2003.
- [12] C. Righi. Ad-nilpotent ideals of a parabolic subalgebra. *J. Algebra*, 319:1555–2584, 2008.
- [13] R. Suter. Abelian ideals in a Borel subalgebra of a complex simple lie algebra. *Inventiones mathematicae*, 156, 2004.

T. RIDENOUR: DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY
E-mail address: tbr4@math.northwestern.edu

A. SANDLER: COLUMBIA UNIVERSITY
E-mail address: ars2240@columbia.edu